

Algebra Examinations of Governorates 2012

Cairo

1

Mathematics department - Ahmed Lotfy (E.LS)

1 Complete each of the following:

- a) The solution set of the inequality $-x > 3$ in \mathbb{R} is
- b) $\sqrt[3]{125} - \sqrt[3]{24} = \dots\dots\dots$
- c) The slope of the straight line parallel to x - axis is
- d) The additive inverse for $(1 - \sqrt{2})$ is
- e) $\sqrt[3]{-64} + \sqrt{16} = \dots\dots\dots$

2 Choose the correct answer:

- a) If $x = 2 + \sqrt{5}$, y is the conjugate number for x then $(x - y)^2 = \dots\dots\dots$ ($2\sqrt{8}$ or 20 or $4\sqrt{5}$ or 10).
- b) $(\sqrt{8} + \sqrt{2})^2 = \dots\dots\dots$ ($\sqrt{10}$ or 10 or 18 or $\sqrt{18}$).
- c) $2 \in \dots\dots\dots$ ($|2, 5|$ or $|2, 5|$ or $\{1, 5\}$ or $|1, 5|$).
- d) The radius length of a right circular cylinder whose volume is $40\pi \text{ cm}^3$ and height 10 cm = cm. ($5, 3, 2, 1$)
- e) The irrational number lies between 2 and 3 is ($\sqrt{10}$ or $\sqrt{7}$ or 2.5 or $\sqrt{3}$)

3 (a) Find the solution set of the inequality: $1 \leq 2x + 3 < 5$ in \mathbb{R} and represent it on the number line.

(b) Put in the simplest form: $\frac{1}{4}\sqrt{80} - \sqrt{20} - \sqrt{25} + \sqrt{125}$

- 4 (a)** Find the slope of the straight line that passes by the ordered pairs $(0, -3)$, $(2, 1)$
- (b) Find the total area of a cuboid whose volume is 750 cm and its height 5 cm. with a squared shape base.
- 5** The following table represent the daily wages by L.E for 30 workers. Find the arithmetic mean of that distribution:

Set.	4 -	8 -	12 -	16 -	Total
Frequency	3	5	14	8	30

Cairo

2

El Khalifa & Mokattam Educational Zone
Futures Languages Schools

Answer the following questions:

1 Choose the correct answer:

1. $Q \cup Q \setminus = \dots\dots\dots$ a) \emptyset b) \mathbb{R} c) Q d) $\{O\}$

- 2) $\sqrt[3]{64} + \sqrt{16} = \dots\dots$
 a) zero b) 8 c) -8 d) ± 8
- 3) $[-2, 5] \cup]4, 6[= \dots\dots\dots$
 a) $[-2, 6[$ b) $] -2, 6[$ c) $[-2, 6[$ d) $] -2, 6[$
4. If $(a, 2a)$ satisfied the equation: $2x + 3y = 24$, then $a = \dots\dots\dots$
 a) 2 b) -3 c) -2 d) 3
5. The mean of the values 4, 7, 12, 13, 8, 10 is $\dots\dots\dots$
 a) 6 b) 8 c) 9 d) 12

2 Complete each of the following:

- $[4, 7] \sim \{4, 7\} = \dots\dots\dots$
- 4 cm, 7 cm and 5 cm are dimensions of cuboid, then its volume = $\dots\dots\dots$
- The S.S of the equation: $x^3 + 9 = 1$ in R is $\dots\dots\dots$
- If the lower limit of a set is 10 and the upper limit of the same set is 20, then its center = $\dots\dots\dots$
- If the ordered pair $(-1, 1)$ satisfies the relation $-x + by = 16$, then $b = \dots\dots\dots$

- 3 (a)** If $x = 2\sqrt{2} - \sqrt{3}$ $y = \frac{5}{2\sqrt{2} - \sqrt{3}}$, find the value of the expression: $(x + y)^2$
 (b) Find in R the S.S of: $8x - 17 \geq 6x + 11$

- 4 (a) By using the number line:** If $x = [-7, 3]$, $y = [-3, \infty[$
 Then Find: 1) $x \cap y$ 2) $x - y$

(b) Find the slope of the straight line \overleftrightarrow{AB} where $A(-1, 3)$, $B(2, 5)$. Is the point $(8, 1) \in \overleftrightarrow{AB}$?

- 5 (a)** The following table shows the frequency distribution of marks of 40 students in math:

Sets	5 -	15 -	25 -	35 -	45 -	Total
Frequency	7	9	12	8	4	40

Find the mean of this distribution.

- b) Find the length of the radius, of the right circular cylinder if its volume $64\pi \text{ cm}^3$ and its height 4 cm. Represent these data by broken line.

Answer the following questions:

2 Complete each of the following:

- The conjugate number of the number $\sqrt{7} - 2 = \dots\dots\dots$
- The median of a set 3, 7, 9, 5 and 4 is $\dots\dots\dots$
- The S.S of the inequality: $-x > 3$ in R is $\dots\dots\dots$ as interval.

d) $[3,7] - \{3,7\} = \dots\dots\dots$

e) If the straight line represents $x-y = 2$ cuts $x - axis$,then $x = \dots\dots\dots$

2 Choose the correct answer:

1) The multiplicative inverse of $\frac{\sqrt{3}}{6}$ is $\dots\dots\dots$

- (a) $\sqrt{3}$ b) $2\sqrt{3}$ c) $3\sqrt{3}$ d) $3\sqrt{6}$

2) $\sqrt{3} \dots\dots\dots [1, 2]$

- (a) \in b) \notin c) \subset d) $\not\subset$

3) The S.S of the equation : $x^2 + 25 = 0$ in R is $\dots\dots\dots$

- (a) 5 b) ± 5 c) ϕ d) -5

4) $\sqrt{2} + \sqrt{8} = \dots\dots\dots$

- (a) $\sqrt{10}$ b) 4 c) $4\sqrt{2}$ d) $3\sqrt{2}$

5. The mode of values 2, 5, 8, 2, 3 is $\dots\dots\dots$

- (a) 4 b) 2 c) 3 d) 8

3 a) Simplify to the simplest form: $\sqrt[3]{16} - \sqrt[3]{54} - \sqrt[3]{128}$

(b) Find the value of k where (k, 2k) satisfies the , relation $x + y = 15$

4 a) Find the S.S in R of the following inequality:

$-1 \leq 2x + 3 < 5$, then represent the S.S. on the number line.

(b) Find the lateral area for a right cylinder of volume

924 cm^3 . and of a height 6 cm ($\pi = \frac{22}{7}$)

5 In the following table find the mean of marks of 50 students in an examination:

Sets	2-	6-	10-	14-	18-	22-	26-	Total
Frequency	3	6	8	10	11	8	4	50

1 Answer the following questions:

- The volume of the sphere = $\dots\dots\dots$
- $\sqrt{64} - \sqrt[3]{64} = \dots\dots\dots$
- The S.S in R for the equation: $x^2 + 4 = 0$ is $\dots\dots\dots$
- $] -2, 1 [\cap [-2, 1] = \dots\dots\dots$
- $(\sqrt{3} - \sqrt{2})^2 + 2\sqrt{6} = \dots\dots\dots$

2 Choose the correct answer from the given ones:

1. The multiplicative inverse of $\sqrt{\frac{3}{6}}$ is $\dots\dots\dots$

- a) $\sqrt{3}$ b) $2\sqrt{3}$ c) $3\sqrt{3}$ d) $3\sqrt{6}$

- 2) If the ordered pair $(-1, 3)$ satisfies the relation $3x - y = C$, then $C = \dots\dots$
 a) 7 b) -7 c) 6 d) -6
- 3) The S.S. in R for the equation $x^3 = -8$ is $\{\dots\dots\}$
 a) 2 b) 4 c) -2 d) -4
4. The cube whose volume is 8 cm^3 , the area of one of its faces = $\dots\dots \text{ cm}^2$
 a) 4 b) 8 c) 16 d) 64
5. If The mode of the values: 4, a, 5, 3 is 3, then $a = \dots\dots$
 a) 4 b) 3 c) 5 d) 6

3 (a) Find the value of k such that \overleftrightarrow{AB} is parallel to y-axis where
 $A(6, 2)$ $B(-2k, 4)$

(b) If $A =]-\infty, 3[$, $B = [-1, 5]$

Find the following using the number line. (1) $A \cap B$ (2) $A \cup B$

4 (a) Simplify to the simplest form:

$$\sqrt{50} - \sqrt{18} - \sqrt{2}$$

(b) $-7 \leq 4x - 3 < 5$, then represent it on the number line.

5 The following is the frequency distribution of the weekly extra wages of 100 workers in a factory:

Extra wages	30-	40-	50-	60-	70-	80-	Total
No. of workers	10	K	20	28	20	8	100

(1) Find the value of k.

(2) Find the mean of this distribution.

Cairo

5

Rod El-Farag directory - Fatma El-Zahraa
 Experimental language School

Answer the following questions:

1 Choose the correct answer from between brackets:

1. $\sqrt{12} - \sqrt{3} = \dots\dots\dots$ (3 or $\sqrt{3}$ or $2\sqrt{3}$ or $3\sqrt{3}$)

2. $[-2, 3] - \{-2, 3\} = \dots\dots\dots$ ($[-2, 3[$ or $]-2, 3]$ or $[-1, 2]$ or $]-2, 3[$)

3. The ordered pair $(3, 1)$ satisfies the relation $\dots\dots\dots$

($x - y = 6$, $3x + y = 6$, $x - 3y = 6$, $x + 3y = 6$)

4. The volume of the sphere whose diameter length is 6 cm = $\dots\dots\dots \text{ cm}^3$

(9π or 2π or 36π or 288π)

5. The multiplicative inverse of the number $\frac{\sqrt{3}}{2}$ is $\dots\dots\dots$

($\frac{3}{2}\sqrt{2}$ or $\frac{3}{2}\sqrt{2}$ or $\sqrt{\frac{6}{3}}$ or $-\sqrt{\frac{3}{2}}$)

2 Complete the following:

1. If $\sqrt[3]{x} = 3$, then $x = \dots\dots\dots$
2. The S.S of the inequality $-3x + 2 \geq 11$ is $\dots\dots\dots$
3. The cube hose lateral area is 36 m^2 , its volume = $\dots\dots\dots$
4. If $x < \sqrt{15} < x + 1$, $x \in \mathbb{Z}$, then $x = \dots\dots\dots$
5. If $(k, 2k)$ satisfied the equation $2x + 3y = 24$, then $k = \dots\dots$

3 (a) If the straight line that represents the relation $y - x = a$ cuts $y -$ axis at $(k, 3)$.

Find the value of a .

(b) Find the S.S of $(2x - 1)^3 - 10 = 54$

4 (a) If $x = 2\sqrt{2} - \sqrt{3}$ and $y = \frac{5}{2\sqrt{2} - \sqrt{3}}$, then calculate $\frac{x+y}{xy}$

(b) Simplify: $\sqrt[3]{54} - 2\sqrt[3]{\frac{1}{4}} - \sqrt[3]{16}$

5 From the following frequency table.

Sets	10-	20-	30-	40-	X-	60-	Total
Frequency	12	15	25	27	$K + 4$	4	100

Find 1) the values of both X and K .

2) The arithmetic mean of this distribution.

Cairo

6

**El Nozha Educational Zone, El Sayeda Khadiga
Language School**

1 Complete:

- 1) The multiplicative inverse of the number $\frac{\sqrt{2}}{12}$ is = $\dots\dots\dots$
2. $[0, 2] \cup [1, 5[= \dots\dots\dots$
3. If $x = \sqrt{3} + \sqrt{2}$, then $x^{-1} = \dots\dots\dots$
4. The arithmetic means of 3, 2, 7, 9, 9 is $\dots\dots\dots$
5. The mode of 8, 2, 1, 8, 1, 8 is = $\dots\dots\dots$

2 Choose the correct answer:

1. If the volume of a sphere is $\frac{32}{3} \pi \text{ cm}^3$, then its radius length = $\dots\dots\dots$ cm
{1, 2, 4, 16}
2. $\sqrt[3]{2} + \sqrt[3]{2} = \dots\dots\dots$ { $\sqrt[3]{3}$, $\sqrt[3]{4}$, $\sqrt[3]{8}$, $\sqrt[3]{16}$ }
3. If A, B and C are collinear, then the slope of $\overleftrightarrow{BC} = \dots\dots\dots$
4. The median of 9, 6, 8, 7, 2 is $\dots\dots\dots$ { 6, 7, 8, 9 }
5. The irrational number between 2, 3 is $\dots\dots\dots$ { $\sqrt{10}$, $\sqrt{7}$, $\sqrt{15}$, $\sqrt{3}$ }

3 a) Put in the simplest form: $\sqrt{48} - 2\sqrt{27} - 6\sqrt{\frac{1}{3}}$

b) Find the lateral area of a right circular cylinder whose diameter length is 4 cm and its height = 10 cm

4 a) Find the S.S of equations:

(i) $2 - \sqrt{6} x = 8$ (ii) $(x + 2)^3 = -8$

(b) Find the S.S of inequality: $1 \leq 1 - 2x < |-7|$

5 a) Find the slope of the straight line \overleftrightarrow{AB} where A(5, 0) , (0, 2).

(b) The following table shows the frequency of marks of 50 student:

Sets	2-	6-	10-	14-	18-	22-	26-	Total
Freq.	3	6	8	10	11	8	4	50

Find the Arithmetic mean of the marks of the students.

Cairo

7

**Rod El Farag Educational Directorate
Ahd Gedid Language School**

1 Choose the correct answer:

1. $\sqrt{12} - \sqrt{3} = \dots\dots\dots$

- (a) 3 b) $\sqrt{3}$ c) $2\sqrt{3}$ d) $3\sqrt{3}$

2. $] -1, 2 [- [1, 4] = \dots\dots\dots$

- (a) $] -1, 1 [$ b) $\{-1, 1\}$ c) $] -1, 1]$ d) $[-1, 1]$

3. The multiplicative inverse of $\sqrt{\frac{3}{6}}$ is $\dots\dots\dots$

- (a) $-\frac{\sqrt{3}}{6}$ b) $6\sqrt{3}$ c) $2\sqrt{3}$ d) $-2\sqrt{3}$

4. The median of a set of the values 3, 7, 2, 9, 5 is $\dots\dots\dots$

- (a) 6 b) 5 c) 7 d) 3

5. The mean of 5, 6, 2, 7 is $\dots\dots\dots$

- (a) 4 b) 5 c) 7 d) 6

2 Complete:

a) The ordered pairs that satisfy the relation $2x - 3y = 6$ are (0, $\dots\dots\dots$) , ($\dots\dots\dots$, 0)

b) $[3, 5] - \{3, 5\} = \dots\dots\dots$

c) The S.S of $x^2 + 4 = 0$ is $\dots\dots\dots$

d) $\sqrt[3]{64} - \sqrt{16} = \dots\dots\dots$

e) The conjugate of $2 - \sqrt{5}$ is $\dots\dots\dots$

3 (a) Simplify : $\sqrt{32} - \sqrt{72} - 6\sqrt{\frac{1}{2}}$

(b) Find the volume of cylinder with base raduis 7 cm and its height 10 cm $\pi = \frac{22}{7}$

4 a) If $x = \sqrt{5} + 2$, $Y = \sqrt{5} - 2$. Find the value of $(x + y)^2$

b) Represent the following relation graphically $x - y = 1$

5 a) Find the S.S of inequality of $-1 < x + 3 \leq 5$ interval

b) Find the mean of the following date.

Sets	10-	20-	30-	40-	50-	Total
Frequency	10	20	25	30	15	100

Cairo

8

El-Zeitoun Directorate - Language School

1 Answer the following questions:

1. The slope of the straight line is undefined when it is parallel to

2. $\sqrt{64} - \sqrt[3]{64} = \dots\dots\dots$

3. The S.S in R for the equation: $x^2 + 4 = 0$ is

4. $] -2, 1 [\cap [-2, 1] = \dots\dots$

5. $(\sqrt{3} - \sqrt{2})^2 + 2\sqrt{6} \dots\dots\dots$

2 Choose the correct answer from the given ones:

1. The multiplicative inverse of $\sqrt{\frac{3}{6}}$ is

- (a) $\sqrt{3}$ b) $2\sqrt{3}$ c) $3\sqrt{3}$ d) $3\sqrt{6}$

2. The S.S in R of the inequality $-x < 0$ is

- (a) $] -\infty, 0]$ b) $] -\infty, 0 [$ c) $] 0, \infty [$ d) $[0, \infty [$

3. The S.S in R for the equation $x^3 = -8$ is {

- (a) 2 b) 4 c) -2 d) -4

4. The cube whose volume is 8 cm^3 , the area of one of it's faces = cm^2

- (a) 4 b) 8 c) 16 d) 64

5. If the mode of the values: 4, a, 5, 3, is 3, then a =

- (a) 4 b) 3 c) 5 d) 6

3 (a) If $x = \sqrt{5} + \sqrt{3}$, $y = \frac{2}{\sqrt{5} + \sqrt{3}}$. Find the value of the expression: $(x - y)^2$

(b) If $A =] -\infty, 3 [$, $B = [-1, 5]$. Find the following using the number line

- (1) $A \cap B$ (2) $A \cup B$

- 4 (a) Find three ordered pairs satisfy the relation $6x = 1 - y$
 (b) $-7 \leq 4x - 3 < 5$, then represent it on the number line.
- 5 The following is the frequency distribution of the weekly extra wages of 100 workers in a factory:

Extra wages	30-	40-	50-	60-	70-	80-	Total
No. of workers	10	k	20	28	20	8	100

- 1) Find the value of k.
 2) Find the mean of this distribution.

Cairo

9

Al Salam Education Zone - Anwer Al Sadat Exp. Lang. School

Answer the following questions:

1 Choose the correct answer:

- If $\frac{3}{a+2}$ is a rational number, then $a \neq$ (3 , 5 , -2 , zero)
- The mean of the values 7, 15, 19, 14 and 15 is (14 , 15, 16, 17)
- The slope of the constant straight line is
- The multiplicative inverse of $\frac{\sqrt{3}}{6}$ is ($-\frac{\sqrt{3}}{6}$, $6\sqrt{3}$, $2\sqrt{3}$, $-2\sqrt{3}$)
- The mode of the values 3 , 5, 3 , 6, 3 and 8 is = (3 , 5 , 6, 8)

2 Complete:

- The solution set of the equation: $x^2 + 4 = 0$ in R is
- The multiplicative neutral in R is
- $[-1, 5] -]-1, 5[=$
- $\sqrt[3]{-8} =$
- If $x < \sqrt{15} < x + 1$, $x \in \mathbb{Z}$, then $x =$

3 a) Find the solution set of the inequality:

$x - 3 \geq 4$ in R and represent it on the number line.

b) Simplify: $\sqrt{32} - \sqrt{72} + 6\sqrt{\frac{1}{2}}$

4 a) Find the volume of cylinder with base radius length 7 cm. and its height 10 cm.
 ($\pi = \frac{22}{7}$)

b) If the straight line $\overleftrightarrow{AB} \parallel x$ -axis where

A (1, 8) , B (-7, 4 K). Find k.

5 a) Find the mean of the following data:

Sets	8-	12-	16-	20-	24-	Total
Frequency	4	10	16	12	8	50

b) Simplify : $(\sqrt{5} - \sqrt{2})^2 + \sqrt{40}$

Giza

10

Omrania Zone - El-Sadat E.L.S

Answer the following questions:

1 Choose the correct answer:

1) If the volume of cube is 64 cm^3 , then a lateral of = cm^2 . (4, 8, 64, 96)

2) $\sqrt[3]{54} + \sqrt[3]{-2} = \dots\dots\dots$ ($\sqrt[3]{52}$, $\sqrt[3]{2}$, $2\sqrt[3]{2}$, $4\sqrt[3]{2}$)

3) The relation $y = 3$ is represented by a straight line cutting the y-axis at

((3, 0), (-3, 0), (0, 3), (0, -3))

4) If the mode of 7, $x + 2$, 5 is 7, then $x = \dots\dots\dots$ (7, 5, 2, 12)

5) $3 \in \dots\dots\dots$ ($]3, 5[$, $]0, 3[$, $\{1, 5\}$, $]1, 5[$)

2 Complete each of the following:

1) $] -3, 2[\cap \mathbb{R}^- = \dots\dots\dots$

2) If $\sqrt[3]{x} = \sqrt{9}$, then $x \dots\dots\dots$

3) The S.S of the $-X > 3$ in \mathbb{R} is

4) The median of 3, 7, 2, 5 and 4 is

5) The volume of the sphere whos radius length $\sqrt[3]{21}$ is

3 (a) If $x = 2\sqrt{2} - \sqrt{3}$ $y = \frac{5}{2\sqrt{2} - \sqrt{3}}$. Find the value of $(x + y)^2$

(b) Find in S.S of the inequality and represent on number line $7x - 12 \geq 5x - 8$.

4 (a) Simplify $= 2\sqrt{18} + \sqrt{50} - \frac{1}{3}\sqrt{162}$

(b) If $A =] -1, 3]$ and $B = [0, 5 [$.

Find each the following using the number line: $A \cap B$ $A - B$.

5 (a) Find the slope of the straight line \overleftrightarrow{AB} where

$A(-6, 7)$, $B(0, 2)$

(b) The following table shows the frequency distribution of marks of 100 students in math:

Sets	10-	20-	30-	40-	50-	Total
Frequency	10	20	25	30	15	100

Find the mean of frequency table.

Giza

11

Omrania Educational Directorate El-Neel Language School

1 Complete:

- (a) The solution set of inequality $-3 \leq -x < 3$ in \mathbb{R} is
- (b) $[-1, 3] - \{3\} = \dots\dots\dots$
- (c) The sphere whose volume is $36\pi \text{ cm}^3$ its radius length = cm
- (d) If the lower limit of a set is 6 and the upper limit of the same set is 10, then its centre =
- (e) The conjugate number of the number $\frac{2}{\sqrt{5} - \sqrt{3}}$ is

2 Choose:

- a) $\sqrt[3]{54} + \sqrt[3]{2} = \dots\dots\dots$ ($3\sqrt[3]{2}, 2\sqrt[3]{2}, 3, 2$)
- b) $2, 5 [\cup] \cup \{2, 5\} = \dots\dots\dots$ ($\emptyset, \{2, 5\}, [2, 5], (2, 5)$)
- c) The real numbers $\mathbb{R} = \dots\dots\dots$ ($\mathbb{R}_+ \cup \mathbb{R}, \mathbb{Q} \cap \mathbb{Q}',]-\infty, \infty [\cup \{0\}$)
- d) $(\sqrt{2} + \sqrt{3})^2 - 2\sqrt{6} = \dots\dots\dots$ ($\mathbb{R}_+ \cup \mathbb{R}, \mathbb{Q} \cap \mathbb{Q}', [\infty, \infty [, \mathbb{R}_+ \cup \{0\}$)
- e) The slope of the straight line $y = 1$ is
- ($-1, 0, 1, \text{undefined}$)

- 3 a) If $x = \sqrt{3} + \sqrt{2}$, and $y = \frac{1}{\sqrt{3} + \sqrt{2}}$, find the value of $x^2 + 2xy + y^2$

b) Simplify: $\sqrt[3]{16} - \sqrt[3]{54} - \sqrt[3]{128} + \sqrt[3]{\frac{1}{4}}$

- 4 (a) Find in \mathbb{R} , the S.S. of the inequality $4x - 3 \geq -7$, then represent it on the number line.

(b) Find the slope of the straight line \overleftrightarrow{CD} where $C(0, -3), D(2, 5)$

- 5 (a) The following is the frequency distribution of the weekly extra wages of 100 workers in a factory.

Extra wages	30-	40-	50-	60-	70-	80-	Total
No. of workers	10	14	20	28	20	8	100

Find the mean of this distribution.

1 Complete:

- $[-2, 6] \cap [2, 7] = \dots\dots\dots$
- The additive inverse of $2 + \sqrt{3}$ is $\dots\dots\dots$
- If $x < \sqrt{15} < x + 1$, $x \in \mathbb{Z}$, then $x = \dots\dots\dots$
- The mean of the value 1, 4, 5, 10 is $\dots\dots\dots$
- The slope of the straight line $x = 2$ is $\dots\dots\dots$

2 Choose the correct answer:

- A sphere of radius 1 cm, then its volume is $\dots\dots\dots \text{cm}^3$.
 a) $\frac{\pi}{2}$ b) $\frac{\pi}{6}$ c) $\frac{4\pi}{3}$ d) $\frac{3\pi}{4}$
- The value of the number $\frac{2}{\sqrt{7} - \sqrt{5}}$ is $\dots\dots\dots$
 a) $\sqrt{7} - \sqrt{5}$ b) $\sqrt{7} + \sqrt{5}$ c) $2\sqrt{7}$ d) $\sqrt{5} - \sqrt{7}$
- The multiplicative inverse of $\frac{\sqrt{2}}{10}$ is $\dots\dots\dots$
 a) $2\sqrt{2}$ b) $5\sqrt{2}$ c) $\sqrt{2}$ d) $\sqrt{10}$
- $[2, 7] - \{2, 7\} = \dots\dots\dots$
 a) $\{2, 7\}$ b) ϕ c) $]2, 7]$ d) $]2, 7[$
- $\sqrt[3]{3} + \sqrt[3]{3} = \dots\dots\dots$
 a) 3 b) $2\sqrt[3]{3}$ c) $\sqrt[3]{9}$ d) $\sqrt[3]{6}$

3 Simplify: If the straight line

$y = 2x + 3$ cuts

x - axis at A and y - axis at B

Find the area of ΔABO where O is the origin.

4 Find the S.S in R of the following:

- $3x - 1 \geq 8$
- $(2x - 1)^3 = 8$

5 The following table shows the weekly wages in pounds of 50 workers in a factory:

Sets	5-	15-	25-	35-	45-	Total
Frequency	7	10	12	13	8	50

Find the mean of the wage of the worker in pounds.

1 Choose the correct answer from the given ones:

(a) $\frac{1}{2} \sqrt{20} = \dots\dots\dots$ $[\sqrt{5}, \sqrt{10}, 2\sqrt{5}, 5\sqrt{2}]$

(b) The cube whose volume is 250 cm^3 , the area of one face of this cube = $\dots\dots\dots$
 cm^2 $[50, 125, 20\sqrt[3]{2}, 25\sqrt[3]{4}]$

(c) $[-\infty, 2] \cup [2, \infty[= \dots\dots\dots$

$(\mathbb{R}, \mathbb{R} - \{-2, 2\}, \mathbb{R} - [-2, 2], \mathbb{R} -]-2, 2[)$

(d) The next number to $\sqrt{2}, \sqrt{8}, \sqrt{18}, \sqrt{32}$ is $\dots\dots\dots$

$[\sqrt{40}, \sqrt{42}, \sqrt{50}, \sqrt{38}]$

(e) The mean of the values 24, 25, 22, 23, 29 = $\dots\dots\dots$ $[23.24, 24.4, 24.6]$

2 Complete the following:

(a) If $x \in [-2, 3] =$, then $2x + 1 \in \dots\dots\dots$ $[\dots\dots\dots, \dots\dots\dots]$

(b) If the slope of \overrightarrow{AB} is equal to the slope of \overrightarrow{BC} then A, B and C are $\dots\dots\dots$

(c) The multiplicative inverse of $\frac{\sqrt{3}}{6}$ is $\dots\dots\dots$

(d) If $x = (\sqrt{5} + \sqrt{3})$, $y = (\sqrt{5} - \sqrt{3})$, then $(xy)^3 = \dots\dots\dots$

(e) The mode of the values 23, 32, 33, 22, 24, 32, 23, 24, 33, 22, 32 is $\dots\dots\dots$

3 (a) Put in the simplest form:

$$\frac{1}{4} \sqrt{80} - \sqrt{20} - \sqrt{45} + \sqrt{125}$$

(b) a- If $x = \sqrt{7} + \sqrt{3}$, $y = \frac{2}{\sqrt{7} + \sqrt{3}}$, prove that x and y are conjugates, then
 find the values of $x^2 - 2xy + y^2$

4 (a) Find the solution set of the inequality $7x - 12 \geq 5x - 8$, in \mathbb{R} , in a form of an interval.

And represent it on the number line.

(b) Graph the relation $x - 2y = 1$, then find the slope of straight line

5 (a) Find the radius of a right circular cylinder whose volume is 40π , and its height is 10 cm?

(b) The following table shows the frequency distribution of the marks of 100 students in the exam. Find the mean of the student's marks?

Sets	10-	20-	30-	40-	50-	Total
Frequency	10	20	25	30	15	100

Menofia

14

Educational Shebeen Elkom Directorate - Idel School

Answer the following questions:

1 Choose the correct answer:

1) $[2, 6] - \{2, 8\} = \dots\dots\dots$

- a) \emptyset b) $\{2,6\}$ c) $[2,6]$ d) $]2,6]$

2) $\sqrt{\dots\dots\dots} = \sqrt[3]{64} \dots\dots\dots$

- a) 4 b) 16 c) 64 d) - 64

3) The ordered pair (0, -3) dose not satisfy the relation

- a) $2x + 3y = 12$ b) $2x - 3y = 12$ c) $3x - 4y = 12$ d) $3x + 4y = 12$

4) The S.S of the equation $x^2 - 9 = 0$ ine R is

- a) $\{3\}$ b) $\{-3\}$ c) $\{\pm 3\}$ d) \emptyset

5) The multiplicative inverse of the number $\frac{2}{\sqrt{3}}$ is

- a) $2\sqrt{3}$ b) $3\sqrt{2}$ c) $-3\sqrt{2}$ d) $\frac{\sqrt{3}}{2}$

2 Complete:

(a) $[3, \infty] \cap]-1, 8] = \dots\dots\dots$

(b) $\sqrt[3]{3} + \sqrt[3]{24} - 3\sqrt[3]{81} = \dots\dots\dots$

(c) $\sqrt{8} \times \sqrt{2} = \sqrt[3]{\dots\dots\dots}$

(d) If the lateral area of a cube is 100 cm^2 . then its volume =

(e) The number $\sqrt{29}$ is included between the two consecutive integers and

3 (a) If $x = \sqrt{7} + \sqrt{3}$, $y = \frac{4}{\sqrt{7} + \sqrt{3}}$. Find the value of $x^2 + 2xy + y^2$

(b) Find the S.S of the following inequality $2x + 8 > 3x - 5 > 2x - 8$, in R in a form of an interval , then represent the S.S on number line.

4 (a) If A(2, 1) , B (3, - 1) , C (0, 5) Are A, B , C collinear.

(b) Find the volume of the cylinder whose radius length is 21 cm and height 4 cm, then find its lateral surface area. ($\pi = \frac{22}{7}$)

- 5 From the following frequency distribution draw the histogram and from the draw find the mode.

Sets	10-	20-	30-	40-	50-	60-	Total
Frequency	15	15	20	25	15	10	100

Draw a bar chart for the frequency table data.

Gharbia 15

Samannoud Educational Directorate Samannoud Experimental Language School

Answer the following questions:

- 1 Choose the correct answer from the given ones:

1) The irrational number lies between 2 and 3.

- a) 4 b) $2\frac{1}{2}$ c) $\sqrt{5}$ d) $\sqrt{10}$

2) $[3, 7] - \{3, 7\} = \dots\dots\dots$

- a) $\{0\}$ b) \emptyset c) $[3, 7]$ d) $]3, 7[$

3) The slope of straight line parallel to y– axis is

4) $\sqrt{12} - \sqrt{3} = \dots\dots\dots$

- a) $\sqrt{9}$ b) $\sqrt{3}$ c) 3 d) 4

5) The mode of the values: 4, 5, 6, 4 and 7 is

- a) 4 b) 5 c) 6 d) 7

- 2 Complete each of the following:

a) If: $-x > \text{zero}$, then S.S in R is

b) The radius length of a sphere whose volume is $\frac{9}{2} \pi \text{ cm}^3 = \dots\dots\dots \text{ cm}$

c) The multiplicative inverse of the number $\sqrt{5} - 2$ is

d) The arithmetic mean of the values: 4, 6, 8 and 10 is

e) The median of the values 5, 10, 8, 12 and 11 is

- 3 (a) Find the slope of \overrightarrow{AB} where A (7, 2), B (-2, 0)

(b) If $A = \sqrt{5} - \sqrt{3}$, $B = \frac{2}{A}$. Find the value of the expression: $A^2 - 2AB + B^2$

- 4 (a) Find in R the S.S of each of the following

- 1) $3x + 5 > 2$ 2) $x^2 + 3 = 19$

(b) Calculate the radius length of the right cylinder whose volume is 2156 cm^3 and its height is 14 cm. ($\pi = \frac{22}{7}$)

5 (a) If $x = [-3, 2[$ and $y = [-1, 4]$, find in the form of interval using the number line each of the following:

- 1) $X \cap Y$ 2) $X - Y$

(b) The following table shows the frequency distribution of marks of 60 students in an exam .

Sets	5-	10-	15-	20-	25-	30-	35-
Frequency	2	5	15	20	13	5	1

Form the ascending cumulative frequency table and represent it graphically.

1 Complete:

- a) $\sqrt[3]{64} - \sqrt{16} = \dots\dots\dots$
 b) The ordered pair (3 ,) satisfies the relation $x - 3y = 9$
 c) $[-1, 5] \cap [3, 7] = \dots\dots\dots$
 d) The median of 23 , 42 , 17 , 30 and 20 is
 e) If the lower limit of a set is 4 and its upper limit is 8 then its centre is

2 Choose the correct answer:

- a) The conjugate of the number $\sqrt{5} - \sqrt{2}$ is
 ($\sqrt{5} + \sqrt{2}$, $\sqrt{5} - \sqrt{2}$, $2\sqrt{5}$, $\sqrt{2} - \sqrt{5}$)
 b) The multiplicative inverse of $\frac{\sqrt{3}}{6}$ is ($6\sqrt{3}$, $2\sqrt{3}$, $\sqrt{6}$, $\sqrt{3}$)
 c) The volume of the cube of side length 3 cm is
 (3 cm^3 , 27 cm^3 , 27 cm , 9 cm)
 d) The mean of 4 , 5 , 6 , 8 , 2 is ($4, 5, 6, 7$)
 e) The S.S. of the equation $x^3 = 64$ is where $X \in R(4, 8, \{4\}, \{8\})$

3 a) If $a = \sqrt{3} + \sqrt{2}$, $b = \frac{1}{\sqrt{3} + \sqrt{2}}$ find the value of $a + b$
in the simplest form.

b) If A (-5 , 6) , B (1 , 3) , C (3 , y) are collinear, find the value of y.

4 a) Find in R. The S.S. of the equation $\sqrt{2}x - 1 = 5$

b) Find the volume of right cylinder whose height is 6 cm and length of its base radius is 1.4 cm
 ($\pi = \frac{22}{7}$)

5 a) Find S.S. in R of the inequality $-1 < 2x + 3 \leq 7$ and represent it on the number line.

b)

Sets	2-	6-	10-	14-	18-	22-	26-
Frequency	3	5	8	10	7	5	2

Using ascending cumulative frequency table to get the median.

Damietta

17

Experimental Language School

Answer the following questions:

1 Choose the correct answer:

a) The relation $10 + y = 5x$ is represented by a straight line cutting x - axis in at

$((2, 0), (-2, 0), (0, 2), (0, -2))$

b) The irrational number between 3 and 4 is $(3.5, \frac{1}{8}, \sqrt{13}, \sqrt{20})$

c) The mean of marks of 5 pupils is 30 then the total of their marks is

$(6, 25, 35, 150)$

d) The S.S. $-x > 3$ in R is

$(\{3\}, |3, |, |-, |3|, |-, -, -3|)$

e) The S.S. $\sqrt{2} X = 2$ in R is

$(\sqrt{2}, 2\sqrt{2}, 2, \emptyset)$

2 Complete to form a correct statement:

a) $Q \cup \bar{Q} = \dots\dots\dots$

b) If $\sqrt[3]{X} = -\sqrt{4}$ then $X = \dots\dots\dots$

c) Complete using \in or \notin $103 \times 10^{-5} \dots\dots\dots R_+$

d) The median 23, 13, 15, 30, 11 is

e) If the lower limit of set is 8 and the upper limit of the same set is 14 then the center is

3 a) Prove that $2\sqrt{32} - \sqrt{50} - \frac{1}{3}\sqrt{162} = 0$

b) If the slope of the straight line passing by $(-1, 3), (x, 2)$ is undefined find x.

4 a) Find the S.S and represent on the number line $15 \leq 5X < 30$ where $X \in R$.

b) If $A = \sqrt{3} + \sqrt{2}$, $B = \frac{1}{\sqrt{3} + \sqrt{2}}$ find value of $A^2 B^2$.

5 a) If $X = [0, 3], Y =] 0, 7]$ find the following using number line.

a) $X \cap Y$

b) $X \cup Y$

b) The following table shows frequency distribution for the scores of 40 students in an exam.

Set	30-	40-	50-	60-	70-	80-	Total
Frequency	3	4	12	8	7	5	40

Graph the frequency histogram, then find the mode.

Answer the following questions:

1 Choose the correct answer:

- 1) The irrational number which lies between 2 and 3 is
 - a) $\sqrt{10}$
 - b) $\sqrt{7}$
 - c) 2.5
 - d) $\sqrt{3}$
- 2) $-5 \dots\dots\dots]-$, $-6[$
 - a) \in
 - b) \notin
 - c) \subset
 - d) $\not\subset$
- 3) $1 + \sqrt{2} \dots\dots\dots \sqrt{3}$
 - a) $<$
 - b) $>$
 - c) $=$
 - d) \geq
- 4) If the ordered pair $(-1, 3)$ satisfies the relation $3X - y = C$ Then $C = \dots\dots\dots$
- 5) The surface area of a square whose side length is $\sqrt{3} = \dots\dots\dots \text{cm}^2$.
 - a) $4\sqrt{3}$
 - b) 9
 - c) 3
 - d) 6

2 Complete:

- 1) The intersection point of the ascending and descending cumulative frequency curves determines on the sets axis.
- 2) $\sqrt[3]{a^3} = \dots\dots\dots$
- 3) The lower limit of a set is 4 and its center is 9 , then its upper limit is =
- 4) If the ages of 5 students are 13 , 15 , 16 , 14 and 17 years old, then the mean of their ages = years.
- 5) $]3, 9[\cup \{9\} = \dots\dots\dots$

3 a) If $x = \frac{4}{\sqrt{7} - \sqrt{3}}$, $y = \sqrt{7} - \sqrt{3}$

Prove that: x and y are conjugates, then find the values of:

- a) $(x + y)$
- b) $x y$

b) If the slope of the straight line \overleftrightarrow{AB} is 7 where $A(1, 2)$, $B(3, x)$ find x.

4 a) If $x = [-2, 3]$, $y = [1, 5[$

Find the following using the number line

- a) $x \cup y$
- b) $x \cap y$

b) Find the solution set for the inequality $3x - 1 < 5$, in \mathbb{R} in the form of interval, then graph the solution on the number line.

5 a) The volume of a sphere is $562.5 \pi \text{cm}^3$. Find its surface area.

b) The following table shows frequency distribution of the daily wages of some workers.

Sets	5-	10-	15-	20-	25-	30-	Total
Frequency	10	14	24	30	12	10	100

Required: Graph the descending cumulative frequency curve.

Answer the following questions:

1 Choose the correct answer:

- 1) The median of the numbers 3, 6, 2, 4, 9 is
 (a) 2 b) 6 c) 4 d) 9
- 2) The mode of the values 3, 5, 3, 6, 3 is
 (a) 3 b) 5 c) 6 d) 8
- 3) Which of the the following ordered pairs satisfies the relation $2x + y = 5$
 (a) (-1,3) b) (1,3) c) (3,1) d) (2,2)
- 4) The additive inverse of $(7 - \sqrt{2})$ is
 (a) $(7 + \sqrt{2})$ b) $(-7 - \sqrt{2})$ c) $(\sqrt{2} - 7)$ d) $\sqrt{5}$
- 5) The S.S of the equation: $x^2 + 16 = 0$ in R is
 (a) ± 4 b) -4 c) 4 d) \emptyset

2 Complete:

- 1) $\mathbb{R}_+ \cap [-1, 3]$
- 2) $\sqrt[3]{64} - \sqrt{16} = \dots\dots\dots$
- 3) The S.S of the inequality $-x > 3$ in R is
- 4) $[1, 7] - \{1, 7\} = \dots\dots\dots$
- 5) If the volume of a cube is 27 cm^3 , then the length of its side is

3 (a) If $x = \sqrt{5} + -\sqrt{2}$, $y = \sqrt{2} - \sqrt{2}$, then find the value of: $\frac{x+y}{xy-1}$

(b) If the slope of the straight line \overleftrightarrow{AB} in $\frac{5}{3}$ where A (3,5), B (-3,4) find the value of y

Answer the following questions:

1 Complete:

- a) If $\sqrt[3]{x} = 5$ then $x = \dots\dots\dots$
- b) $(k,2)$ satisfies the relation $6x + 5y = 13$ then $k = \dots\dots\dots$
- c) The conjugate of $5 - \sqrt{7}$ is

d) The S.S of $x^2 + 25 = 0$ in \mathbb{R} is

e) If the lower limit of a set is 4 and the upper limit of the same set is 10 then the centre of the set =

2 Choose the correct answer:

a) Q' ($3, \sqrt{3}, \sqrt{16}, \sqrt[3]{8}$)

b) $\{x: x \in \mathbb{R}, x > 0\}$ ($\mathbb{R}^+, \mathbb{R}^-, \mathbb{R}^*, \mathbb{R}$)

c) The multiplicative inverse of $\frac{\sqrt{3}}{6}$ is ($-\frac{\sqrt{3}}{6}, 6\sqrt{3}, 2\sqrt{3}, -2\sqrt{3}$)

d) If the mode of the values 3, 5, $x + 1$, 5, 3, 1 is 5 then $x =$ ($5, 4, 3, 6$)

e) If the edge length of a cube is 4 cm then the volume of a cube = cm^3 ($4, 16, 64, 12$)

3 (a) Find in \mathbb{R} the S.S of:

$4 < 3x + 4 \leq 7$ and represent it on the number line.

(b) If:

$A = \sqrt{3} + \sqrt{2}$, $B = \frac{1}{\sqrt{3} + \sqrt{2}}$ Find the value of $A^2 + B^2$

4 (a) Find the slope of the straight line that passes through the two ordered pairs (2,0), (0,3)

(b) If:

$X =] - \infty, 4]$, $Y =] - 2, 7]$

Find each of the following using the number line:

1) $X \cap Y$ 2) $X \cup Y$ 3) $X - Y$

5 (a) Find the volume of a right circular cylinder its radius length 7 cm and its height is 10 cm. ($\pi = \frac{22}{7}$)

(b) Find the mode of the following table:

Sets	2 -	6 -	10 -	14 -	18 -	22 -
Frequency	3	5	6	10	7	2

Answer the following questions:

1 Choose the correct answer:

- 1) $Q \cap Q'$
 - (a) $\{0\}$
 - (b) \emptyset
 - (c) R
 - (d) Q
- 2) The mean of the value 7, 15, 19, 14 and 15
 - (a) 14
 - (b) 15
 - (c) 16
 - (d) 17
- 3) If the volume of a sphere $\frac{32}{3} \pi \text{ cm}^3$, then its radius length = cm
 - (a) 14
 - (b) 2
 - (c) 4
 - (d) 16
- 4) $[-1, 3] - \{3\} =$
 - (a) $] -1, 3 [$
 - (b) $[-1, 3]$
 - (c) $] -1, 3]$
 - (d) $[-1, 3 [$
- 5) The S.S of $x^2 = 16 = 0$, $x \in Q$ is
 - (a) 4
 - (b) ± 4
 - (c) 16
 - (d) 8

2 Complete:

- 1) The mode of the numbers 6, 10, 8, 4, 6, 7, 6 equals
- 2) The additive inverse for $(1 - \sqrt{2})$ is
- 3) The slope of the straight line $y = 2$ is
- 4) $] - \infty, 2] \cup [-3, \infty [=$
- 5) The solution set in R of the inequality $2x + 3 > 13$ is

3 (a) Put in the simplest form:

$$\sqrt{18} + 3\sqrt{\frac{1}{3}} - \sqrt{8} - \sqrt[3]{9}$$

(b) Find the solution set in R $(2x + 3)^3 + 4 = 12$

4 (a) A right circular cylinder of height 10 cm. and its volume is 1540 cm^3 . Find its total area.

(b) Find in R the S.S. of the inequality $5x + 7 \geq 2x - 11$

5 (a) Find the value of k where (k, k) satisfies the relation $2x + y = 12$

(b) Find the mean of the following table:

Sets	5 –	15–	25–	35–	45 –	Total
Frequency	7	9	12	8	4	40

1 Choose the correct answer:

- 1) $4 \in \dots\dots\dots$
 (a) $] -3 .\infty[$ b) $] -\infty .3 [$ c) $(3, 5)$ d) $] 4 .\infty[$
- 2) The slope of the straight line $x = 1$ in $\dots\dots\dots$
 (a) -1 b) 0 c) 1 d) undefined
- 3) The mean of the numbers: 7, 8, 3 and 6 is $\dots\dots\dots$
 (a) 5 b) 6.5 c) 6 d) 8
- 4) If the volume of a sphere = $36 \pi \text{ cm}^3$ then the radius is $\dots\dots\dots$ cm.
 (a) 4 b) 27 c) -3 d) 3
- 5) The multiplicative inverse of $\frac{\sqrt{2}}{6}$ is $\dots\dots\dots$
 (a) $\sqrt{3}$ b) $\sqrt{6}$ c) $3\sqrt{2}$ d) $-\frac{3}{\sqrt{2}}$

2 Complete:

- 1) $\sqrt[3]{-64} + \sqrt{16} = \dots\dots\dots$
- 2) The mode of the values: 5, 3, 8, 5 and 9 is $\dots\dots\dots$
- 3) The S.S in R of the equation: $x^2 - 5 = 0$ is $\dots\dots\dots$
- 4) The conjugate of the number $\frac{2}{\sqrt{5} - \sqrt{3}}$ in the simplest form is $\dots\dots\dots$
- 5) The median of the values: 3, 9, 2, 7, 6 and 11 is $\dots\dots\dots$

3 (a) If $X =] -1 .4]$, $Y = [2. \infty [$ Find: 1) $X \cap Y$ 2) $X \cup Y$

- (b) Find the solution set of the following inequality in R $-3 \leq 2x - 1 < 5$ then represent the S.S. on the number line.**

4 (a) Put in the simplest form:

$$\sqrt{32} - \sqrt{72} + 6\sqrt{\frac{1}{2}}$$

- (b) Find the volume of the right circular cylinder of radius length is 7 cm. and its height is 10 cm. ($\pi = \frac{22}{7}$)**

5 (a) If A (-1,2), B (3,0), C (k,-1) are on the same straight line find the value of k

- (b) The following table shows the frequency distribution of weight of 40 children in k.g:**

Sets	5 –	15–	25–	35–	45 –	Total
Frequency	7	9	12	8	4	40

Calculate the mean of weight of children