

**The First
Note book
First prep**

**The Rational
Numbers**

**Prepared by
Mr /Ahmed Omar**

Set of Rational Numbers

Remember:-

(1) Counting Numbers = $\{1, 2, 3, 4, \dots\}$

(2) Set of Integers $\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$

(3) Set of Natural Number = $\{0, 1, 2, 3, \dots\}$

(4) $\mathbb{Z}^+ = \{1, 2, 3, \dots\}$

(5) $\mathbb{Z}^- = \{-1, -2, -3, \dots\}$

(6) $\mathbb{N} \subset \mathbb{Z}$

(7) $\mathbb{Z} = \mathbb{Z}^+ \cup \{0\} \cup \mathbb{Z}^-$

(8) If $|x| = a$ then $x = \pm a$

Ex:1:- Complete

(1) $\mathbb{Z} - \mathbb{N} = \dots$

(2) $\mathbb{Z}^+ \cup \{0\} = \dots$

(3) $\mathbb{Z}^+ \cap \mathbb{Z}^- = \dots$

(4) $\mathbb{N} \cap \mathbb{Z} = \dots$

(5) $|-3| = \dots$

(6) $|0| = \dots$

(7) If $|x| = 7$ then $x = \dots$ or $x = \dots$

(8) If $|x| = 0$ then $x = \dots$

Ex:2, find the value of x when

(1) $x + 2 = 1 - 3$

(2) $|x| + 2 = 5$

(3) $|x| - 5 = 2 + 1 - 2$

(4) $2x + 5 = 7$

Definition of the rational number:

It's a number that can be expressed in the form

$\frac{a}{b}$ where a and b are integers and $b \neq 0$

i.e.

$$\mathbb{Q} = \{x: x = \frac{a}{b}, a \in \mathbb{Z}, b \in \mathbb{Z}, b \neq 0\}$$

Ex: put \in or \notin

1) $\frac{2}{3} \dots \mathbb{Q} \in$ the answer Because $\frac{2}{3}$ in the form $\frac{a}{b}, b \neq 0$

2) $0.2 \dots \mathbb{Q} \in$ $0.2 = \frac{2}{10}$ in the form $\frac{a}{b}, b \neq 0$

3) $25\% \dots \mathbb{Q} \in$ $25\% = \frac{25}{100}$ in the form $\frac{a}{b}$

4) $1\frac{1}{4} \dots \mathbb{Q} \in$ $1\frac{1}{4} = \frac{5}{4}$ in the form $\frac{a}{b}$

5) $-2 \dots \mathbb{Q} \in$ $-2 = -\frac{2}{1}$

6) zero $\dots \mathbb{Q} \in$ zero = $\frac{0}{1}$

7) $3 \dots \mathbb{Q} \in$ $3 = \frac{3}{1}$

8) $\frac{1}{0} \dots \mathbb{Q} \notin$ because the denominator = 0

9) $\frac{2}{5-5} \dots \mathbb{Q} \notin$ because the denominator = 0

Notes

1) $\frac{a}{b} \in \mathbb{Q}$ if $b \neq 0$ (denominator \neq zero)

2) $\frac{a}{b} = 0$ if $a = \text{zero}$, $b \neq \text{zero}$

3) $\frac{a}{b}$ is an integer if the numerator is divisible by the denominator.

~~5~~

ii) $\frac{x-3}{4}$ since $\frac{x-3}{4} = 0$ therefore $x-3=0$
i.e. $x=3$

Since $\frac{2-x}{3} = 0$ therefore $2-x=0$ i.e. $x=2$

③ $\frac{x+3}{x+4}$

Since $\frac{x+3}{x+4} = 0$ therefore $x+3=0$ i.e. $x=-3$

④) $\frac{3x-3}{x}$

Since $\frac{3x-3}{x} = 0$ therefore $3x-3=0$
 $\Rightarrow 3x=3 \div 3$ i.e. $x=1$

(5) $\frac{2x}{x+3}$

Since $\frac{2x}{x+3} = 0$ therefore $2x = 0 \div 2 \Rightarrow x = 0$

(6) $\frac{3x+6}{x+1}$

Since $\frac{3x+6}{x+1} = 0$ therefore

$$3x + 6 = 0$$
$$3x = -6 \quad \div 3$$

$$x = -2$$

Different forms of a rational number

1 Writing the rational number in its simplest form:

Ex: put each of the following numbers in its simplest form:

$$\textcircled{1} \frac{10}{15} = \frac{2}{3}$$

($\div 5$) Dividing the two terms by 5

$$\textcircled{2} \frac{45}{20} = \frac{9}{4}$$

($\div 5$)

$$\textcircled{3} \left| -\frac{18}{12} \right| = \frac{18}{12} = \frac{3}{2}$$

Dividing the two terms by 6

$$\textcircled{4} \left| 20\% \right| = \frac{20}{100} = \frac{1}{5}$$

Dividing the two terms by 20

$$\textcircled{5} -\frac{24}{28} = -\frac{6}{7}$$

Dividing the two terms by 4

* the form of a terminating decimal

To write the rational number in the form of a terminating decimal we make its denominator equal to 10, 100, 1000, ...

Ex: write each of the following numbers in the form of a terminating decimal:

$$\textcircled{1} \frac{2}{5}$$


(To make denominator 10 multiplying the two terms by 2)

$$\frac{2}{5} = \frac{4}{10} = 0.4$$

Comparing and ordering Rational Numbers.

Ex:1 Represent the rational number $\frac{3}{4}$ on the number line

Solution:-




Ex:2: Represent the rational numbers $2, 0, \frac{3}{2}, -\frac{5}{2}, -3$ on the number line then arrange them descendingly

Solution:

We should convert their denominators to have common denominator at first

Since L.C.M of the denominators is 2
Then the numbers after converting their denominators are

$$\frac{4}{2}, \frac{0}{2}, \frac{3}{2}, -\frac{5}{2}, -\frac{6}{2}$$



Then: $\frac{4}{2} > \frac{3}{2} > \frac{0}{2} > -\frac{5}{2} > -\frac{6}{2}$

i.e $2 > \frac{3}{2} > 0 > -\frac{5}{2} > -3$

Ex3 Compare between the two rational numbers

$$\frac{2}{3} \text{ and } \frac{3}{4}$$

Solution

We should convert their denominators to have common denominator at first

$$\frac{2}{3} \xrightarrow{(1)} \frac{8}{12} \quad \frac{3}{4} \xrightarrow{(2)} \frac{9}{12} \Rightarrow \frac{8}{12} \quad \frac{9}{12}$$

Since $\frac{8}{12} < \frac{9}{12}$ then $\frac{2}{3} < \frac{3}{4}$

Ex4: find four rational numbers lying between the two numbers $\frac{1}{2}$ and $\frac{1}{3}$

Solution

$$\text{Q1 } -\frac{3}{5} \text{ and } -\frac{2}{3}$$

$$\textcircled{1} \quad \frac{1}{2} \xrightarrow{(1)} \frac{3}{6} \quad \frac{1}{3} \xrightarrow{(2)} \frac{2}{6} \Rightarrow \frac{3}{6} \text{ and } \frac{2}{6}$$

$$\Rightarrow \frac{30}{60} \text{ and } \frac{20}{60}$$

$$\Rightarrow \frac{20}{60} < \frac{21}{60} < \frac{22}{60} < \frac{23}{60} < \frac{24}{60} < \frac{30}{60}$$

$$\textcircled{2} \quad -\frac{3}{5} \xrightarrow{(1)} -\frac{9}{15} \quad -\frac{2}{3} \xrightarrow{(2)} -\frac{10}{15} \Rightarrow -\frac{9}{15} \text{ and } -\frac{10}{15}$$

$$\Rightarrow -\frac{90}{150} \text{ and } -\frac{100}{150}$$

then the numbers are: $-\frac{91}{100}, -\frac{92}{100}, -\frac{93}{100}, -\frac{94}{100}$

Ex:5 Complete each of the following using the suitable Sign ($<$, $>$ or $=$):

① $\frac{7}{5} \dots \frac{4}{5}$

② $-\frac{3}{4} \dots -\frac{2}{4}$

$$\boxed{3} \frac{1}{5} \dots \frac{1}{6}$$

④ $\frac{3}{6} \dots \frac{2}{3}$

⑤ $\frac{4}{10} \dots \frac{14}{35}$

6) $1 - \frac{10}{15} \dots \frac{2}{3}$

Try by Yourself

① Represent on the number line each of the following rational numbers

① $\frac{1}{3}$ ② $-\frac{7}{4}$ ③ $-\frac{1}{3}$ ④ $1\frac{1}{5}$ ⑤ $-3\frac{1}{2}$

2) Compare between

① $\frac{2}{3}$ and $\frac{5}{7}$ ② $\frac{1}{5}$ and $\frac{1}{6}$ ③ $-\frac{8}{15}$ and $-\frac{2}{3}$

Q4) 0.6 and $\frac{5}{6}$

[3] find two rational numbers lying between

① $\frac{4}{5}$ and $\frac{3}{4}$ ② $\frac{4}{5}$ and $\frac{5}{6}$ ③ $\frac{4}{5}$ and 0.7 ④ $\frac{3}{4}$, 2

4] Represent the rational numbers: 2 , $-\frac{5}{2}$, $\frac{7}{2}$, 0 , -1 on the number line then arrange the ascendingly

⑤ Complete each of the following using the suitable sign ($<$, $>$ or $=$):

① $\frac{3}{4} \dots \frac{1}{5}$

② $\frac{-3}{15} \dots \frac{-2}{5}$

$$\boxed{3) \frac{3}{15} \dots \frac{4}{20}}$$

④ $-\frac{6}{9} \dots -\frac{2}{6}$

5) $\frac{3}{2} \dots \frac{3}{5}$

⑥) Complete each of the following

① $\frac{3}{5} > \dots > \frac{2}{5}$

② $-\frac{1}{3} > \dots > -\frac{2}{3}$

3) $\frac{1}{4} > \dots > \frac{1}{8}$

④ $\frac{-3}{14} > \dots > \frac{-2}{7}$

⑤ $\frac{1}{4} > \dots > \frac{1}{6}$

(6) $-\frac{2}{9} > \dots > -\frac{5}{6}$

Mr /Ahmed Omar

Adding and Subtracting Rational Numbers

① Adding two rational numbers having the same denominator:

If $\frac{a}{b}$ and $\frac{c}{b}$ are two rational numbers

then:

$$\frac{a}{b} + \frac{c}{b} = \frac{a+c}{b}$$

② Adding two rational numbers with different denominators:

If $\frac{a}{b}$ and $\frac{c}{d}$ are two rational numbers

then:

$$\frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{bd}$$

Example ① Add:

$$① \quad \frac{1}{5} + \frac{2}{5} = \frac{1+2}{5} = \frac{3}{5}$$

$$② \quad \frac{5}{6} + \left(-\frac{1}{6}\right) = \frac{5-1}{6} = \frac{4}{6} = \frac{2}{3} \quad (\div 2)$$

$$③ \quad \frac{7}{8} + \left(-\frac{5}{8}\right) = \frac{7-5}{8} = \frac{2}{8} = \frac{1}{4} \quad (\div 2)$$

$$④ \quad -\frac{1}{4} + \frac{3}{4} = \frac{-1+3}{4} = \frac{2}{4} = \frac{1}{2} \quad (\div 2)$$

$$⑤ \quad \frac{5}{9} + \left(-\frac{2}{9}\right) = \frac{5-2}{9} = \frac{3}{9} = \frac{1}{3} \quad (\div 3)$$

$$⑥ \quad -\frac{2}{3} + \left(-\frac{1}{3}\right) = \frac{-2-1}{3} = \frac{-3}{3} = -1$$

Ex: Write the addition property which used in each of the following

$$\textcircled{1} \frac{7}{2} + \frac{9}{16} = \frac{9}{16} + \frac{7}{2}$$

$$\textcircled{2} \text{Zero} + \frac{3}{4} = \frac{3}{4}$$

$$\textcircled{3} \left(\frac{2}{3} + \frac{1}{3} \right) + \frac{1}{6} = \frac{2}{3} + \left(\frac{1}{3} + \frac{1}{6} \right)$$

$$\textcircled{4} \frac{3}{4} + \left(-\frac{3}{4} \right) = \text{Zero}$$

$$\textcircled{5} \frac{5}{8} + \text{Zero} = \frac{5}{8}$$

Second: Subtraction operation:

If $\frac{a}{b}$ and $\frac{c}{d}$ are two rational numbers, then

$$\frac{a}{b} - \frac{c}{d} = \frac{a}{b} + \left(-\frac{c}{d} \right)$$

i.e. The subtraction operation in \mathbb{Q} is defined as adding the minuend $\left(\frac{a}{b} \right)$ to the additive inverse of the subtrahend $\left(\frac{c}{d} \right)$

① $a - b$

Solution

$$(2) (a+b) - c$$

$$= \left(\frac{3}{4} + \frac{-5}{2} \right) - \frac{1}{2}$$

$$= \left(\frac{6-20}{8} \right) - 12$$

$$= -\frac{14}{8} - \frac{1}{2} = -\frac{7}{4} - \frac{1}{2} = \frac{-14-4}{8} = \frac{-18}{8} = \frac{-9}{4}$$

Multiplying and Dividing Rational Numbers

First: Multiplication operation

If $\frac{a}{b}$ and $\frac{c}{d}$ are two rational numbers

$$\text{then: } \frac{a}{b} \times \frac{c}{d} = \frac{a \times c}{b \times d}$$

Remember:

$$+ \times + = +, \quad - \times - = +$$

$$+ \times - = -, \quad - \times + = -$$

Ex: Find the result of each of the following in its simplest form:

$$\textcircled{1} \quad \frac{3}{6} \times \frac{2}{5} = \frac{3 \times 2}{6 \times 5} = \frac{6}{30} = \frac{1}{5}$$

$$\textcircled{2} \quad -\frac{3}{4} \times \frac{2}{9} = \frac{-3 \times 2}{4 \times 9} = \frac{-6}{36} = -\frac{1}{6}$$

$$\textcircled{3} \quad \frac{1}{2} \times (-2) = \frac{1 \times -2}{2 \times 1} = \frac{-2}{2} = -1$$

$$\textcircled{4} \quad -\frac{7}{8} \times \frac{4}{14} = \frac{-7 \times 4}{8 \times 14} = \frac{-1 \times 1}{2 \times 2} = -\frac{1}{4}$$

$$\textcircled{5} \quad 25\% \times (-\frac{3}{4}) = \frac{25}{100} \times (-\frac{3}{4}) = \frac{1}{4} \times -\frac{3}{4} = -\frac{3}{16}$$

$$\textcircled{6} \quad -1\frac{1}{2} \times -0.3 = -\frac{3}{2} \times -\frac{1}{3} = \frac{-3 \times -1}{2 \times 3} = \frac{3}{6} = \frac{1}{2}$$

$$\frac{b}{a} \times \frac{a}{b} = 1$$

the multiplicative inverse property

Ex: Complete

- ① The multiplicative inverse of the number $\frac{3}{4}$ is ...
- ② The multiplicative inverse of the number -5 is ...
- ③ The multiplicative inverse of the number 0.5 is ...
- ④ The multiplicative inverse of the number $2\frac{1}{2}$...
- ⑤ The multiplicative inverse of the number $(-\frac{2}{3})^{\text{zero}}$...
- ⑥ The multiplicative inverse of the number -1 is ...

Remarks

- ① The multiplicative inverse of the rational number is called the reciprocal of the rational number
- ② Zero has no multiplicative inverse
- $\frac{1}{0}$ is meaningless
- ③ multiplying any rational number by zero equals zero

6 The distributive property:

Ex: Use the Distributing property to find the value of each of the following:

$$\underline{\underline{75}} \quad \frac{13}{12} \times 7 + \frac{13}{12} \times 5$$

$$= \frac{13}{12} \times (7+5) = \frac{13}{12} \times \frac{12}{1} = 13$$

$$\textcircled{2} \quad \frac{4}{9} \times \textcircled{11} + \frac{4}{9} \times \textcircled{16}$$

$$= \frac{4}{9} \times (11 + 16) = \frac{4}{9} \times \frac{27}{1} = 12$$

$$\underline{(3)} \quad \frac{5}{13} \times 2 + \frac{5}{13} \times 4 + \frac{5}{3} \times 7$$

$$= \frac{5}{13} \times (2+4+7) = \frac{5}{13} \times 13 = 5$$

$$\boxed{4} \quad \frac{4}{7} \times 4 + \frac{4}{7} \times 2 + \frac{4}{7} \times 1$$

$$= \frac{4}{7} \times (4 + 2 + 1)$$

$$= \frac{4}{7} \times 7 = 4$$

⑤ $-\frac{3}{7} \times 8 + 5 \times (-\frac{3}{7}) + (-\frac{3}{7})$

$$= -\frac{3}{7} \times (8+5+1) = -\frac{3}{7} \times 14 = -6$$

24

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \times \frac{d}{c}$$

$$\textcircled{1} \frac{1}{2} \div \frac{3}{5} = \frac{1}{2} \times \frac{5}{3} = \frac{1 \times 5}{2 \times 3} = \frac{5}{6}$$

$$\textcircled{2} \quad -\frac{3}{5} \div \frac{0}{5} = -\frac{3}{5} \times \frac{5}{0} = \frac{-15}{0} = -\frac{1}{0}$$

$$\textcircled{3} \quad -\frac{14}{15} \div \left(-\frac{21}{5}\right) = -\frac{14}{15} \times -\frac{5}{21} = \frac{2}{3} \times \frac{1}{3} = \frac{2}{9}$$

④ $2\frac{1}{4} \div 1\frac{1}{2} = \frac{9}{4} \div \frac{3}{2} = \frac{9}{4} \times \frac{2}{3} = \frac{18}{12} = \frac{3}{2}$

Ex: find the value of x in each of the following

$$\textcircled{1} \frac{5}{7} \times x = \frac{5}{7}$$

$$\textcircled{2} \quad x \times \frac{17}{3} = 1$$

③ $-\frac{7}{3} \times x = \text{zero}$

$$\textcircled{4} \quad -\frac{7}{3}x - \frac{3}{7} = x$$

$$(5) \frac{3}{5} \times x = -\frac{4}{5} \times \frac{3}{5}$$

$$(8) \frac{3}{4} \div x = \frac{3}{4} \times -\frac{2}{5}$$

Ex: Find a rational number lying at one fourth of the way between $\frac{1}{2}$, $\frac{1}{3}$

Solution

$$\frac{1}{2}, \frac{1}{3} \rightarrow \frac{3}{6}, \frac{2}{6}$$

The distance = $|1\frac{3}{6} - 2\frac{1}{6}| = \frac{1}{6}$

The required number is

the smaller number + by distance

$$= \frac{2}{6} + \frac{1}{4} \times \frac{1}{6}$$

$$= \frac{2}{6} + \frac{1}{24} = \frac{8+1}{24} = \frac{9}{24} = \frac{3}{8}$$

Ex: Find a rational number lying at one fifth of the way between $\frac{2}{5}$ and $\frac{4}{7}$

Solution $\frac{2}{5}, \frac{4}{7} \Rightarrow \frac{14}{35}, \frac{20}{35}$

the distance = $\left| \frac{20}{35} - \frac{14}{35} \right| = \frac{6}{35}$

the required number =

the smaller numbers $\frac{1}{5}$ distance

$$= \frac{14}{35} + \frac{1}{5} \times \frac{6}{35} = \frac{14}{35} + \frac{6}{175} = \frac{70+6}{175} = \frac{76}{175}$$